

**vitaforce**

VITAFORCE RES-Q REMEDY

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS

Unscheduled

PROPRIETARY NAME AND DOSAGE FORM

RES-Q REMEDY Drops

READ ALL OF THIS LEAFLET CAREFULLY BECAUSE IT CONTAINS IMPORTANT INFORMATION FOR YOU
RES-Q REMEDY is available without a doctor's prescription, for you to treat a mild illness. Nevertheless, you still need to use RES-Q REMEDY carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Do not share RES-Q REMEDY with any other person.
- Ask your health care provider or pharmacist if you need more information or advice.

WHAT RES-Q REMEDY CONTAINS

The active substance is:

Composition per 100 g:	
Prunus Cerasifera (Cherry Plum) [Flower]	0.1 g
Clematis Vitalba (Clematis) [Flower]	0.1 g
Helianthemum Nummularium (Rock Rose) [Flower]	0.1 g
Impatiens Glandulifera (Impatiens) [Flower]	0.1 g
Ornithogalum Umbellatum (Star of Bethlehem) [Flower]	0.1 g

The other ingredients are:
Ethanol and purified water.
Contains: Ethanol 22% m/m.

WHAT RES-Q REMEDY IS USED FOR

Category D Complementary Medicine
Discipline 33.6 Western Herbal Remedy
An herbal remedy for shock.

Supports the treatment of stress and anxiety.

BEFORE YOU TAKE RES-Q REMEDY

Do not take RES-Q REMEDY if you are hypersensitive (allergic) to any of the ingredients. This unregistered medicine has not been evaluated by SAHPRA for its quality, safety or intended use.

TAKE SPECIAL CARE WITH RES-Q REMEDY

Disulfiram: The alcohol content of Bach flower remedies can cause a disulfiram reaction when taken within 12 hours of disulfiram. Patients taking disulfiram should not take any supplements containing alcohol including Bach flower remedies. Metronidazole can cause a disulfiram-like reaction when taken with alcohol. Star of Bethlehem contains cardiac glycosides similar to digitoxins. Theoretically, taking Star of Bethlehem with other cardiac glycoside-containing products might increase the risk of cardiac glycoside toxicity. RES-Q REMEDY may make you sleepy.

TAKING RES-Q REMEDY WITH FOOD AND DRINK

RES-Q REMEDY can be taken with food.

PREGNANCY AND BREASTFEEDING

Not safe for use during pregnancy and lactation. This preparation contains alcohol, which is teratogenic and dangerous for foetal development.

DRIVING AND USING MACHINERY

- RES-Q REMEDY may make you feel drowsy/sleepy.
- Do not drive because RES-Q REMEDY could interfere with your ability to drive safely.
- Do not operate any tools or machines.

IMPORTANT INFORMATION ABOUT SOME OF THE INGREDIENTS OF RES-Q REMEDY

Sugar free
Suitable for vegans and vegetarians.
Contains: Ethanol 22% m/m.

TAKING OTHER MEDICINES WITH RES-Q REMEDY

Always tell your healthcare provider if you are taking any other medicine.

HOW TO TAKE RES-Q REMEDY

Do not share medicines prescribed for you with any other

person. Always take RES-Q REMEDY exactly as instructed. You should check with your doctor or pharmacist if you are unsure.

The usual dosage is:

Adults and children from 12 years of age: 3 to 5 drops to be taken in a little water every five minutes for acute cases, until symptoms improve. Thereafter reduce to every two hours.
From 6-12 years of age: 1-3 drops
From 4 months - 2 years of age: 1 drop
Do not exceed the stated dose.
Shake bottle before use.

IF YOU TAKE MORE RES-Q REMEDY THAN YOU SHOULD

In the event of overdosage, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison control centre.

IF YOU FORGET TO TAKE A DOSE OF RES-Q REMEDY

Do not take a double dose to make up for forgotten individual doses.

POSSIBLE SIDE EFFECTS

RES-Q REMEDY can have side effects.

There are no known side effects.

Not all side effects reported for this medicine are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking this medicine, please consult your healthcare provider for advice. If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

STORING AND DISPOSING OF RES-Q REMEDY

Store all medicines out of reach of children.

Store at or below 25 °C.

Keep the container in the outer carton.

Store in the original package / container.

Keep the container tightly closed.

Protect from light.

Protect from moisture.

Do not store in a bathroom.

Do not use after the expiry date stated on the label / carton / bottle.

Do not use RES-Q REMEDY if you notice visible signs of deterioration.

Return all unused medicine to your pharmacist.

Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

PRESENTATION OF RES-Q REMEDY

A carton containing 20 ml or 50 ml amber dropper bottle.

IDENTIFICATION OF RES-Q REMEDY

A clear colourless liquid.

REGISTRATION NUMBER

U4138 Act 101/1965

DATE OF REGISTRATION

TBA

PROFESSIONAL INFORMATION

For further information refer to the Professional Information of the product.

NAME AND ADDRESS OF REGISTRATION HOLDER

Ascendis Consumer Brands (Pty) Ltd
31 Georgian Crescent East
Bryanston, Gauteng 2191
South Africa
+27 11 036 9420

DATE OF PUBLICATION OF THE PROFESSIONAL INFORMATION

TBA



VITAFORCE RES-Q REMEDY

PASIËNTINLIGTINGSBLAADJIE

SKEDULERINGSTATUS

Ongeskeduleer

HANDELSNAAM EN DOSEERVORM

RES-Q REMEDY Druppels

LEES DIE HELE INLIGTINGSBLAADJIE NOUKEURIG DEUR, DIT BEVAT BELANGRIKE INLIGTING VIR JOU.
RES-Q REMEDY is beskikbaar sonder 'n doktersvoorskrif om minder ernstige siektes te behandel. Jy moet egter nogsteeds RES-Q REMEDY versigtig gebruik om die beste resultate te verkry.

- Hou hierdie voubiljet, dit mag dalk nodig wees om dit weer te lees.
- Moenie RES-Q REMEDY met enige ander persoon deel nie.
- Raadpleeg jou gesondheidsoorgryskyn of apoteker indien jy meer inligting van advies benoeft.

WAT RES-Q REMEDY BEVAT

Die aktiewe bestanddele is:

Samestelling per 100 g :

Waterige ekstrak van die blomme van:	
Prunus Cerasifera (Cherry Plum) [Blom]	0.1 g
Clematis Vitalba (Clematis) [Blom]	0.1 g
Helianthemum Nummularium (Rock Rose) [Blom]	0.1 g
Impatiens Glandulifera (Impatiens) [Blom]	0.1 g
Ornithogalum Umbellatum (Star of Bethlehem) [Blom]	0.1 g

Die ander bestanddele is:

Ethanol en gesuwerde water.

Bevat: Ethanol 22% m/m.

WAARVOOR RES-Q REMEDY GEBRUIK WORD

Kategorie D Komplementêre Medisyne

Discipline 33.6 Westerse Kruie

In Kruie behandeling vir emotionele skok.

Ondersteun die behandeling van spanning en angs.

VOORDAT RES-Q REMEDY GEBRUIK WORD

Moenie RES-Q REMEDY gebruik indien jy hipersensitie (allergie) is vir enige van die bestanddele nie. Hierdie ongeregistreerde medisyne is nie deur SAHPRA geëvalueer vir sy gehalte, veiligheid of beoogde gebruik nie.

NEEM SPESIALE SORG MET RES-Q REMEDY

Disulfiram: Die alkoholinhoud van Bach-blom geneesmiddels kan 'n disulfiram-reaksie veroorsaak as dit binne 12 uur na disulfiram geneem word. Pasiënte wat disulfiram neem, mag geen aanvullings vat wat alkohol bevat nie, insluitende Bach-blom geneesmiddels. Metronidazole kan 'n reaksie soortgelyk aan disulfiram veroorsaak wanneer dit saam met alkohol geneem word. Star of Bethlehem bevat hartglykoside soortgelyk aan digitale. Teoreties mag die glykydine ineen van "Star of Bethlehem" en ander hart glykoside produkte die risiko van hart glykoside vergiftiging verhoog. RES-Q REMEDY mag jou daar slaperig laat voel.

NEEM VAN RES-Q REMEDY MET KOS EN DRANK

RES-Q REMEDY kan saam met kos geene word.

SWANGERSKAP EN BORSVOEDING

Nie veilig tydens swangerskap en borsvoeding. Hierdie samestelling bevat alkohol, wat teratogen is en gevraaglik is vir fetaale ontwikkeling.

Die BESTUUR EN GEBRUIK VAN MASJINERIE

- RES-Q REMEDY mag jou dalk lomerig/slaperig laat voel.
- Moenie bestuur nie want RES-Q REMEDY kan you vermoë te veilig bestuur belemmer.
- Moenie gereedskap of masjiene gebruik nie.

BELANGRIKE INLIGTING IN VERBAND MET SOMMIGE VAN DIE BESTANDDELE IN RES-Q REMEDY

Swikervry

Geskik vir streg vegetariërs.

Bevat: Ethanol 22% m/m.

AS JY ANDER MEDISYNE SAAM MET RES-Q**REMEDY NEEM**

Stel altyd jou gesondheidsorgpraktisyen in kennis indien jy enige ander medisyne gebruik.

HOE OM RES-Q REMEDY TE GEBRUIK

Moenie jou medisyne met enige iemand anders deel nie. Neem RES-Q REMEDY altyd presies soos aanbeveel. Bevestig met jou dokter of apoteker indien jy onseker is. Die normale dosering is:

Drie tot vyf druppels in 'n bietjie water, om geneem te word elke vyf minute in akute gevalle, tottad die simptome verbeter. Kan daarna verminder word tot elke twee ure.

Van 6 tot 12 jaar oud: 1-3 druppels

Van 2 tot 6 jaar oud: 1-2 druppels

Van 4 maande tot 2 jaar oud: 1 druppel

Moet nie die aanbevolle dosis oorskry nie.

Skud die bottel voor gebruik.

INDIEN JY MEER RES-Q REMEDY GEBRUIK AS WAT JY BEHOORT

Indien oordosering vermoed word, raadpleeg onmiddellik jou dokter of apoteker. Indien geenlike beskikbaar is nie, kontak die naaste hospitaal of gibeerhersentrum.

INDIEN JY VERGEET HET OM RES-Q REMEDY TE NEEM

Moenie 'n dubbele dosis neem om vir oorgeslange dosisse te vergoed nie.

MOONTLIKE NEWE-EFFEKTE

RES-Q REMEDY kan moontlik newe-effekte veroorsaak.

Daar is geen newe-effekte bekend nie.

Nie alle newe-effekte wat vir hierdie medisyen aangemeld is word in hierdie voubiljet genoem nie. Indien jou algemene gesondheid verswak of indien jy enige nadelige effekte ondervind terwyl jy hierdie medikasie gebruik, moet jy asseblief jou dokter, apoteker of ander gesondheidsorgpraktisyen raadpleeg. Indien jy enige newe-effekte waarnem wat nie in hierdie voubiljet genoem word nie, moet jy jou dokter of apoteker daarvan vertel.

BERGING EN WEGDOEN VAN RES-Q REMEDY

Bewaar alle medikasie buiten bereik van kinders.

Bewaar teen of benede 25 °C.

Bewaar die houer in die kartonverpakking.

Bewaar in die oorspronklike houer.

Maak die houer dig toe.

Beskerm teen lig.

Beskerm teen vog.

Moenie die produk in die badkamer bewaar nie.

Moenie na die vervaldatoon wat op die etiket / houer / bottel aangedui is, gebruik nie.

Moenie RES-Q REMEDY gebruik indien jy enige sigbare tekenens van produkafbraak opmerk nie.

Besorg enige ongebruikte medisyne terug aan jou apoteker.

Moenie van ongebruikte medisyne ontslae raak in dreine of rioolstelsels nie (bv. toilette).

AANBIEDING VAN RES-Q REMEDY

'n 20 ml of 50 ml amberkleurige botteltjie met 'n drupper in kartonverpakking.

IDENTIFIKASIE VAN RES-Q REMEDY

'n Heldere, kleurlose vloeistof.

REGISTRASIONOMMER

U4138 Act 101/1965

DATUM VAN PUBLIKASIE

TBA

PROFESSIONELE INLIFTING

Verwys na die voubiljet met professionele inliting vir meer inliting.

NAAM EN BESIGHEIDSADRES VAN DIE HOUER VAN DIE REGISTRASIESERTIFIKAAT

Ascendis Consumer Brands (Pty) Ltd

31 Georgian Crescent East

Bryanston, Gauteng 2191

South Africa

+27 11 036 9420

DATUM VAN PUBLIKASIE VAN DIE PROFESSIONELE INLIFTING

TBA

**VITAFORCE
RES-Q REMEDY****PROFESSIONAL INFORMATION****SCHEDULING STATUS**

Unscheduled

PROPRIETARY NAME AND DOSAGE FORM

RES-Q REMEDY Drops

COMPOSITION

Composition per 100 g:

Aqueous infusion of the flowers of:	
Prunus Cerasifera (Cherry Plum) [Flower]	0,1 g
Clematis Vitalba (Clematis) [Flower]	0,1 g
Helianthemum Nummularium (Rock Rose) [Flower]	0,1 g
Impatiens Glandulifera (Impatiens) [Flower]	0,1 g
Ornithogalum Umbellatum (Star of Bethlehem) [Flower]	0,1 g

Inactive ingredients:

Ethanol and purified water.

Contains: Ethanol 22% v/m

Sugar free

Suitable for vegans and vegetarians.

PHARMACOLOGICAL CLASSIFICATION

Category D Complementary Medicine

Discipline 33.6 Western Herbal Remedy

PHARMACOLOGICAL ACTION

RES-Q REMEDY is a flower remedy that was originally discovered by Dr Edward Bach. Dr Bach believed that illness was caused by a negative state of mind. RES-Q REMEDY is manufactured to his original concept, and is specifically formulated to help with shock, stress, anxiety, fainting and other conditions associated with accidents and emergencies.

INDICATIONS

An herbal remedy for shock.

Supports the treatment of stress and anxiety.

CONTRAINDICATIONS

There are no known contraindications.

WARNINGS AND SPECIAL PRECAUTIONS

Do not use this product if you are hypersensitive (allergic) to any of the ingredients. If you are on any medication or suffering from any medical condition, it is advisable to seek medical advice before starting any new medicine, supplement or remedy.

Do not exceed recommended dosage(s) unless on the advice of a healthcare practitioner.

This unregistered medicine has not been evaluated by SAHPRA for its quality, safety or intended use.

INTERACTIONS

Disulfiram : The alcohol content of Bach flower remedies can cause a disulfiram reaction when taken within 12 hours of disulfiram. Patients taking disulfiram should not take any supplements containing alcohol including Bach flower remedies.

Metronidazole can cause a disulfiram-like reaction when taken with alcohol.

Star of Bethlehem contains cardiac glycosides similar to digitoxin. Theoretically, taking Star of Bethlehem with other cardiac glycoside-containing products might increase the risk of cardiac glycoside toxicity.

HUMAN REPRODUCTION

Not safe for use during pregnancy and lactation.

This preparation contains alcohol, which is teratogenic and dangerous for foetal development.

DOSAGE AND DIRECTIONS FOR USE

Adults and children from 12 years of age: 3 to 5 drops to be taken in a little water every five minutes for acute cases, until symptoms improve. Thereafter reduce to every two hours.

From 6-12 years of age: 1-3 drops

From 2-6 years of age: 1-2 drops

From 4 months -2 years of age: 1 drop

Do not exceed the stated dose.

Shake bottle before use.

SIDE EFFECTS

There are no known side effects.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT

Treatment should be symptomatic and supportive. In the case of accidental overdose of the product, contact the nearest hospital or poison control centre.

IDENTIFICATION

A clear colourless liquid.

PRESENTATION

A carton containing 20 ml or 50 ml amber dropper bottle.

STORAGE INSTRUCTIONS

Store at or below 25 °C. Protect from light.
KEEP OUT OF REACH OF CHILDREN.

REGISTRATION NUMBER

U4138 Act 101/1965

DATE OF REGISTRATION

TBA

NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION

Ascendis Consumer Brands (Pty) Ltd

31 Georgian Crescent East

Bryanston, Gauteng 2191

South Africa

+27 11 036 9420

DATE OF PUBLICATION OF THE PROFESSIONAL INFORMATION

TBA

08A0013/01

